*Journal of Immunological Methods*, 160 (1993) 275–276

© 1993 Elsevier Science Publishers B.V. All rights reserved 0022-1759/93/$06.00

# Contents to volume 160 (1993)

**Number 1 – 15 March 1993**

*Research reports*

Comparison between direct binding, competition and agglutination assays in the characterization of polyclonal anti-idiotypes against anti-HBs human monoclonal antibodies
*J.M. Cocchi, M.A. Trabaud, J. Grange, P.F. Serres and C. Desgranges (France)* 1

Development of a two-site immunoradiometric assay for rat/human corticotrophin-releasing factor. Application to the measurement of interleukin-1$\beta$-stimulated production of hypothalamic CRF in vitro
*P. Hagan, F. Tilders, S. Poole and A.F. Bristow (UK, Netherlands)* 11

Generation of human monoclonal antibodies by transformation of lymphoblastoid B cells with ras oncogene
*S. Shammah, T.L. Mantovani, R. Dalla-Favera and P. Casali (USA)* 19

Fluorescent europium chelates as target cell markers in the assessment of natural killer cell cytotoxicity
*K. Blomberg and A.-C. Ulfstedt (Finland)* 27

Selective biotinylation of *Neisseria meningitidis* group B capsular polysaccharide and application in an improved ELISA for the detection of specific antibodies
*J. Diaz Romero and I. Outschoorn (Spain)* 35

Electroporation and commercial liposomes efficiently deliver soluble protein into the MHC class I presentation pathway. Priming in vitro and in vivo for class I-restricted recognition of soluble antigen
*W. Chen, F.R. Carbone and J. McCluskey (Australia)* 49

Graphic analysis of superoxide secretion by phagocytic leukocytes at different cell concentrations
*G.L. Tritsch and R.L. Evans (USA)* 59

A sensitive method to detect defined peptide among those eluted from murine MHC class II molecules
*T.D. Brumeanu, R. Kohanski, C.A. Bona and H. Zaghouani (USA)* 65

Improved rapid photometric assay for quantitative measurement of PMN migration
*W.G. Junger, T.A. Cardoza, F.C. Liu, D.B. Hoyt and R. Goodwin (USA)* 73

The use of ATP bioluminescence as a measure of cell proliferation and cytotoxicity
*S.P.M. Crouch, R. Kozlowski, K.J. Slater and J. Fletcher (UK)* 81

A new approach for measurement of cytotoxicity using colorimetric assay
*R.F. Hussain, A.M.E. Nouri and R.T.D. Oliver (UK)* 89

Assessment of the suitability of commercially available SpA affinity solid phases for the purification of murine monoclonal antibodies at process scale
*M.A.J. Godfrey, P. Kwasowski, R. Clift and V. Marks (UK)* 97

Quantitation of vitronectin and clusterin. Pitfalls and solutions in enzyme immunoassays for adhesive proteins
*K. Høgåsen, T.E. Mollnes, J. Tschopp and M. Harboe (Norway, Switzerland)* 107

An efficient procedure for the generation of human monoclonal antibodies based on activation of human B lymphocytes by a murine thymoma cell line
*J. Kwekkeboom, M. Van Oosten, M. De Boer, L. Van Alphen, M.L.C.M. Mevissen, E. Lindhout, J.M. Tager and C. De Groot (Netherlands, USA, Italy)* 117

*Short communications*

Determination of antibody affinity by ELISA with a non-linear regression program. Evaluation of linearized approximations
*R.W. Glaser (Germany)* 129

Use of stable 6-aminohexyl derivatives for labelling polysaccharides with haptens and for preparing polysaccharide immunoadsorbents
*P.L. Ey (Australia)* 135

*Letter to the editors*

A safe and rapid method for analyzing apoptosis-induced fragmentation of DNA extracted from tissues or cultured cells
*P. Prigent, C. Blanpied, J. Aten and F. Hirsch (France)* 139

*Software notice*
CBEIA: programs for simulation of ELISA experiments and affinity determination
*R.W. Glaser (Germany)* 141

*Announcements* 143

*Notes to authors*

**Number 2 – 2 April 1993**

*Immunological standardization*
The international standard for interleukin-6. Evaluation in an international collaborative study
*R.E. Gaines Das and S. Poole (UK)* 147

*Research reports*
Improvement of antisera raised against complex antigen mixtures by the use of heterologous sources of antigen for immunization
*P. Hammerl, A. Hartl and J. Thalhamer (Austria)* 155
The use of monoclonal antibodies specific for seal immunoglobulins in an enzyme-linked immunosorbent assay to detect canine distemper virus-specific immunoglobulin in seal plasma samples
*D.P. King, A.W.M. Hay, I. Robinson and S.W. Evans (UK)* 163
In vivo and in vitro primed lymphocytes. Correlation of cytochemically detected BLT-specific lymphoid serine protease with cytotoxic activity
*L. Wagner, G. Sunder-Plassmann, W. Base, M. Wiesholzer, V. Sexl, G. Lang and C.P. Worman (Austria, UK)* 173
Cytokine-specific ELISPOT assay. Single cell analysis of IL-2, IL-4 and IL-6 producing cells
*K. Fujihashi, J.R. McGhee, K.W. Beagley, D.T. McPherson, S.A. McPherson, C.-M. Huang and H. Kiyono (USA)* 181
Epitope mapping and binding kinetics of monoclonal antibodies studied by real time biospecific interaction analysis using surface plasmon resonance
*B. Johne, M. Gadnell and K. Hansen (Norway, Sweden)* 191
Preparation of an anti-acid sphingomyelinase monoclonal antibody for the quantitative determination and polypeptide analysis of lysosomal sphingomyelinase in fibroblasts from normal and Niemann-Pick type A patients
*R. Rousson, P. Parvaz, J. Bonnet, C. Rodriguez-Lafrasse, P. Louisot and M.T. Vanier (France)* 199
Comparison of four purification methods for the production of immunoglobulins from eggs laid by hens immunized with an enterotoxigenic *E. coli* strain
*E.M. Akita and S. Nakai (Canada)* 207
Inactivation of HIV-infected H9 cells in whole blood preparations by lysing/fixing reagents used in flow cytometry
*J.K.A. Nicholson, S.W. Browning, S.L. Orloff and J.S. McDougal (USA)* 215
Phenotypic and functional analysis of mucosal T cells isolated from tissue explants of rat upper respiratory tract
*C. McMenamin, B. Chilna and P.G. Holt (Australia)* 219
An algorithm for confirming screen reactivity in blood donors in enzyme immunoassays for antibodies to hepatitis C virus
*J. Craske, W.K. Paver and D. Farmer (UK)* 227
Fractionation of schistosome antigens by high performance electrophoretic chromatography and their screening for the ability to induce Th1 lymphocyte activity
*A.P. Mountford and R.A. Wilson (UK)* 237
Complete saturation of protamine sulphate by dsDNA is necessary in order to obtain a highly sensitive and specific anti-dsDNA ELISA
*A. Rupin, W. De Jong, D. Degenne and P. Bardos (France)* 245
An immunoblotting procedure for screening glycophorins and band 3 protein in the same blots. Identification of glycophorin and band 3 variant forms
*K. Wasniowska, M. Czerwinski, J. Halasa and E. Lisowska (Poland)* 253
An ELISA for blood group specific exoglycosidases
*L. Hobbs, R. Phillips and D. Smith (USA)* 261
Improved bioassay for the detection of porcine tumor necrosis factor using a homologous cell line: PK(15)
*G. Bertoni, P. Kuhnert, E. Peterhans and U. Pauli (Switzerland)* 267

*Book reviews* 273

*Contents to volume 160 (1993)* 275

*Notes to authors* 277